

# Radon

This article is a 'work in progress' incorporating new information whenever time permits.

1. Building materials
2. The effect of changes in building construction techniques
3. Occupational exposures
4. Ground disturbances (earthquakes, etc.)
5. Fracking
6. Weather and seasonal changes
7. Water supplies
8. Health effects
9. Remediation
10. Measuring radon levels in your home, school and workplace
11. 108 References

## Radon

Radon is considered the second cause of lung cancer and the first in those who have never smoked (Torres-Durán [2013](#), Choi & Mazzone [2014](#), Laurier & Gay [2015](#), Choi [2016](#)). Hinojosa de la Garza ([2014](#)) reported that it was evident that lung cancer mortality was directly associated with locations with high levels of radon. Their study took into account a population which had been stable for more than 25 years, which they believe suggested spatial clustering of lung cancer deaths due to indoor radon concentrations. Geologic unit, well water, community, weather and unconventional natural gas development were associated with indoor radon concentrations (Casey [2015](#)). Stopping smoking at age 50 years decreases the lifetime risk due to radon by around a half relative to continuing smoking, but the risk for ex-smokers remains about a factor of 5-7 higher than that for never-smokers.

In an Irish study of people with lung cancer (Smyth [2016](#)), 21 % of patients had radon levels recorded above the national reference level. Only 5 % of patients were aware of the association between radon gas and lung cancer. Smokers were significantly less likely to engage fully in radon testing. In another study from Ireland (Dowdall [2016](#)), it is estimated that approximately 7% of the national housing stock have radon concentrations above the Reference level of 200 Bq m<sup>-3</sup>.

Radon mitigation used to reduce radon concentrations in homes can also have a substantial impact on lung cancer risk, even for persons in their 50's. Smokers in high-radon homes should both stop smoking and remediate their homes (Hunter [2015](#)). In a study of lung cancer in never-smokers, by Torres-Durán ([2015](#)), the median residential radon concentration was 195 Bq m<sup>-3</sup>, higher than the action levels recommended by the World Health Organisation.

Alpha particle emissions from inhaled radon decay products, and not radon itself, cause lung cancer. Alpha particles, which can only penetrate a short distance into bronchial epithelium, induce more biological damage than beta or gamma radiation, and can induce DNA base mutations and chromosomal strand breaks (Melloni [2014](#)).

Radon and its decay products are the cause of 50% of the total radiation dose from natural sources. Thoron gas is one of the most common isotopes of radon. Indoor radon and thoron concentrations are not correlated, though thoron levels are comparatively low, between 0.5 and 6% of the radiation dose.

Public Health England recommends that radon levels should be reduced in homes where the average is more than 200 Bq m<sup>-3</sup>. This recommendation has been endorsed by the Government. The Target Level of 100 Bq m<sup>-3</sup> is the ideal outcome for remediation works in existing buildings and protective measures in new buildings. If the result of a radon assessment is between the Target and Action Levels, action to reduce the level should be seriously considered, especially if there is a smoker or ex-smoker in the home. Axelsson ([2015](#)) suggested that 35-40% of the radon attributed lung cancer cases can be prevented if radon levels higher than 100 Bq m<sup>-3</sup> are lowered to 100 Bq m<sup>-3</sup>.

Worldwide, 3-20 % of all lung cancer deaths are likely caused by indoor radon exposure (SH Kim [2016](#)). Lung cancer deaths induced by radon was slightly higher among females than males. If all homes with radon above 100 Bq/m<sup>3</sup> were effectively remediated reports Kim, studies in Germany and Canada found that 302 and 1704 lung cancer deaths could be prevented each year, respectively. 5 % of all cases of lung cancer in South Korea might be attributable to radon. Studies in Germany, France, and Switzerland reported indoor radon concentrations similar to those in South Korea (Kang [2016](#)).

In Canada, the estimated average thoron concentration is 9 becquerels per metre cubed ( $\text{Bq m}^{-3}$ ), and the average radon concentration is  $96 \text{ Bq m}^{-3}$ , more than double the worldwide indoor radon concentration (Chen [2014](#)). A study in Canada (Rauch & Henderson [2013](#)) concluded “The radon potential map of Canada may communicate potential radon risk, but it was not designed for epidemiologic exposure assessment. Overall, the potential map classified 34 LHAs as higher than observed, and 10 LHAs as lower than observed. The potential map should only be used to inform exposure assessment in conjunction with observed radon concentrations.” The results of a study by Branion-Calles ([2015](#)) contribute evidence supporting the use of a reference level lower than the current guideline of  $200 \text{ Bq m}^{-3}$  for the province of British Columbia.

The Environmental Protection Agency (EPA) in the USA and Public Health England (PHE) estimate that there are 22,000 radon-related deaths in the USA each year and over 1,100 in the UK as a result of inadequate preventative measures and poor detection methodologies. The Environmental Protection Agency (EPA) announced a strategy in 2015 for preventing 3,200 lung cancer deaths annually by 2020, by reducing radon levels in five million homes, schools and childcare centres.

In **Kosovo**, in two villages affected by depleted uranium, radon levels ranged from 82 to  $432 \text{ Bq m}^{-3}$ ; the value of  $400 \text{ Bq m}^{-3}$  was exceeded in 2 of the 25 homes measured (Nafezi [2014](#)). At least 15% of homes in Finland (Valmari [2014](#)) were found to exceed their reference levels of  $400 \text{ Bq m}^{-3}$ .

In the **USA**, the results of the 1992 EPA National Radon Survey estimated that 1 in 15 homes (about 5.8 million) had an elevated radon level (Keith [2012](#)).

The radon levels obtained in most assessed offices in Obafemi Awolowo University in **Nigeria** were found to be within the permissible reference levels. It was recommended that mitigation measures should be put in place in the few offices above permissible levels (Afolabi [2015](#)).

In places of high radon concentration  $^{222}\text{Rn}$  daughters adhere to clothes, skin and hair, adding some radiative concentration to that due to radon and its progeny (Martín Sánchez [2013](#)).

A survey by Nursan ([2014](#)) concluded that there was a necessity to inform the public about significant environmental risks such as radon gas. Well-educated middle-aged parent of high school students seemed to be inadequately informed. Signorelli & Limina ([2002](#)) said that due to the populations’ low risk perception (caused by unawareness of the problem) radon is undoubtedly the environmental pollutant which has the most impact on public health. In Italy 4,000 cases of lung cancer are attributable to radon (about 11% of total lung cancer) have been estimated per year.

A survey of 40 homes, 31 elementary schools and 5 kindergartens in **Macedonia** (Stojanovska [2016](#)) found measured values varied from 22 to  $990 \text{ Bq/m}^3$  for indoor radon concentrations. It seems that the variation in level of radon exposure suggests that it would be worth measuring field levels and taking remedial action where necessary.

**Serbian** radon levels depended on the geological background, with elevated levels in areas with vulcanite and granitoid rocks (Bossev [2013](#)). A study by Zunić ([2014](#)) confirmed the strong influence of geological factors in the **Balkan** region on the variability of indoor radon, and large differences of radon concentrations could be found in different rooms of the same house and significant difference in radon concentrations in one season. Extensive measurements of radon in the town of Xanthi in northern **Greece** show that the part of the town overlying granite deposits and the outcrop of a uranium ore has exceptionally high indoor radon levels, with monthly means up to  $1500 \text{ Bq m}^{-3}$ . 40% of the properties in this part of the town exhibit radon levels above  $200 \text{ Bq m}^{-3}$  while 11% of the houses had radon levels above  $400 \text{ Bq m}^{-3}$  (Kourtidis [2015](#)).

The European Council directive of 5 December 2013 on basic safety standards for protection against the dangers arising from exposure to ionising radiation (2013/59/Euratom) tightened up the radon levels allowed in the workplace and exposures in homes are regulated for the first time (Bochicchio [2014](#)). Barros ([2015](#)) concluded that *“Because of the percentage of workplaces with elevated radon concentrations, additional surveys of workplace radon concentrations are needed, especially in areas of high radon potential, to assess the contribution of workplace radon exposure to an individual’s overall radon exposure.”*

## Building materials

One of the most important indoor pollutants is radon from soils and building materials (Schram-Bijkerk [2013](#), Kumar [2013](#), Borgoni [2014](#), Singh [2016](#)). In a study in Romania, the main source of radon was the building sub-soil and the soil near the house (Cosma 2015). Indoor radon levels are lower in flats than houses, and in buildings with concrete foundations (Kropat [2013](#)).

Radon levels were investigated in homes in **Turkey** where construction materials had included many different sorts of building materials. The levels were between 23 and 1141 Bq m<sup>-3</sup> (Aykamis [2013](#), Kayakökü [2016](#)). Carbonate rock tends to produce higher levels of indoor radon, whereas sedimentary rocks can produce only low levels (Kropat [2013](#)). The Iranian study by Pirsahab concluded that granite stone and adobe coverings could not be recommended for construction purposes.

## The effect of changes in building construction techniques

It can accumulate in residential buildings. Breathing radon and radon progeny for extended periods can be hazardous to health and can lead to lung cancer. Air change rate, indoor temperature and moisture have significant effects on indoor radon concentration. Minimum radon levels were obtained at temperatures between 20 and 22°C and a relative humidity of 50-60% (Akbari [2013](#)). Kropat ([2013](#)) found higher levels of radon at higher temperatures.

Modern energy-efficient architectural solutions and building construction technologies in combination with effective insulation reduce the air permeability of the building. As a result, the air exchange rate is significantly reduced and conditions for increased radon accumulation in indoor air are created. A study by Mohery ([2013](#)), short-lived radon decay products were measured in poorly ventilated living rooms. The maximum radon deposition fractions were found in the upper bronchial airway. In Russia, Yarmoshenko ([2014](#)), found a remarkable increase in the indoor radon concentration level in energy-efficient multi-storey buildings compared with older properties. Zhukovsky & Vasilyev ([2014](#)) also found radon concentration over 200 Bq m<sup>-3</sup> when ventilation rates were low. Vasilyev & Yarmoshenko ([2016](#)) found that in comparison with 2000, as a result of energy-efficient measures in building construction, average radon concentration is likely to increase by a factor of 1.42 by 2030, the percentage above the reference level of 300 Bq/m<sup>3</sup> may increase by a factor of 4.

Radon levels in homes of people with lower socio-economic status are, on average, only about two thirds of those of the more affluent. The reasons for this is not known with certainty, but may be connected with greater underpressure in warmer and better-sealed homes (Kendall [2016](#)).

Jiránek & Kačmařikova ([2014](#)) found that replacing existing windows by new ones decreased the annual energy need for heating 2.8 times, but also reduced the ventilation rate. As a consequence, the 1 year average indoor radon concentration values increased 3.4 times. The additional risk of lung cancer in the thermally retrofitted house increased to a value that is 125 % higher than before conversion.

## Occupational exposures

Radon activity was measured in 34 workplaces in Algiers nuclear research centre between March 2007 and June 2013. The indoor radon levels ranged from 2 to 628 Bq m<sup>-3</sup> (Ait Ziane [2014](#)). With no intervention on occupational radon exposure, estimated lung cancer mortality by age 90 was 16%. Lung cancer mortality was reduced for all interventions (Edwards [2014](#)).

Duan ([2015](#)) found a dose-response association between occupational environmental radon exposure and the risk of lung cancer. The increased risk is particularly apparent when the cumulative exposure to radon is beyond that resulting from exposure to the recommended limit concentration for a prolonged period of time.

In a study by Rossetti & Esposito ([2015](#)), as a result of 8695 measurements in underground workplaces, it was found that the mean radon concentration is higher than that from previous maps elaborated for dwellings and a significant radon concentration was also found in regions traditionally considered as low-risk areas. Elevated levels of radon in public workplaces in Brisbane, Australia were largely found in basements and ground floor levels and in rooms with concrete flooring (Alharbi & Akber [2015](#)).

In Castañar cave, in Spain, cave guides received a total effective dose of 6.41 mSv in 4 months. Public visitors would receive about 12% of the total effective dose permitted at each visit in autumn and about 8-9% in summer (Alvarez-Gallego [2015](#)).

In Polish caves which are open to tourists, as well as providing jobs, it was found that in 67% of the routes the average concentration of radon exceeded 300 Bq m<sup>-3</sup> and in 22 underground routes it exceeded 1000 Bq m<sup>-3</sup> (Olszewski [2015](#)). This may be an organizational, legal and health problem. It is necessary to develop a program of measures to reduce radon concentrations in underground routes, especially routes located in the former mines.

Radon exposure was determined in the Saqqara region in Cairo (El-Kameesy [2016](#)), in tombs and pyramids. Solutions were proposed to protect the health of those who work inside the tombs for long periods of time.

A Chinese study of 3 different kinds of mines were monitored throughout a year (D Fan [2016](#)). The results showed that the individual exposure to radon significantly varied with types of mines and work. Compared with the exposure to coal miners, the exposure to copper miners was much higher. The results indicate that the individual monitoring of radon is necessary for an accurate assessment of radon exposure to miners.

Eleven cohorts of miners occupationally exposed to relatively high concentrations of radon showed a statistically significantly high risk of lung cancer, in a Korean study by Oh ([2016](#)). The authors recommended that preventive policies are needed in order to actively reduce radon exposure and lung cancer incidence in nonsmokers.

Radon activity concentrations (in water and in air) were measured at 13 selected locations at the Avalon Springs thermal spa resort in Montagu (Western Cape, South Africa) to estimate the associated effective dose received by employees and visitors (Botha [2016](#)). The most significant potential radiation exposure identified is that due to inhalation of air rich in radon and its progeny by the resort employees.

## Ground disturbances (earthquakes, etc.)

Tectonic movement and meteorological events are accompanied by radon release. Not only magnitude of earthquakes but also distance from the measurement site should be used for

identifying radon anomalies (Içhedef [2014](#), Briestensky [2014](#), Zhou [2015](#)). Residents in Fukushima were concerned about radon exposure as well as other sources of radiation following the Nuclear accident at Fukushima in 2011 (Tamari [2016](#)).

Areas of high radon activity were located near areas of high flow accumulation showing new submarine groundwater discharge (Rapaglia [2015](#)).

Geologic unit, well water, community, weather, and unconventional natural gas development were associated with indoor radon concentrations (Casey [2015](#)).


Large local radon anomalies may be an indicator of a clandestine underground nuclear explosion (Carrigan [2016](#)).

## **Fracking**

Domestic gas supplied by the controversial process of fracking will increase levels of radioactive radon in people's homes, public health experts have warned. A risk assessment by Public Health England (PHE) shows that shale gas would result in individual exposures to radon that are 15 times higher than through existing supplies of natural gas.

A US health study has found that the hydraulic fracking of unconventional rock formations can liberate and accelerate the release of radon, a highly carcinogenic gas (Casey [2015](#)). US homes in Pennsylvania have been on the rise ever since fracking of the Marcellus shale began in 2004. Buildings in areas of the most active shale gas mining had significantly higher readings of radon compared to buildings located in areas of low well density and fracking activity. Averaged over the whole study period, houses and other buildings using well water had a 21 percent higher concentration of radon than those using municipal water. Houses and buildings located in rural and suburban townships, where most of the gas wells are, had a 39 percent higher concentration of radon than those in cities. In the past decade buildings have been more tightly sealed, potentially trapping radon that gets inside and leading to increased indoor radon levels.

The US findings collaborate and strengthen earlier studies by Australian researchers at Southern Cross University as well as recent shale gas research in Colorado. In [2013](#) Tait reported that the radon air levels above the heavily fracked Surat Basin in Queensland, Australia were three times greater than those observed in a non-fracking region. (A separate and earlier study also showed that methane and carbon dioxide levels in the air were three times higher in the mined landscape.) See model below.



Shallow fracking of coal seams radically changed "the geological structure" of the ground by cracking rock, removing water and lowering pressure in the formation. The fracturing of the ground structure simply created more pathways for radon and other gases to leak up through the soil.

The mechanism for releasing radon into the atmosphere in fracked landscapes may be similar to that caused by earthquakes, added the researchers. Atmospheric levels of radon will typically increase fivefold prior to an earthquake due to stress changes in rock which, in turn, opens new fractures and pathways to the surface.

In 2012 Health Canada found that more than seven per cent of the homes measured in Calgary, Peace River, David Thompson and Aspen regional health districts had dangerous levels of radon (200 becquerels per cubic metre). High levels of radon have also been found in northeast B.C. That geographic region has also witnessed intense shale gas activity over the last decade.

## Weather and seasonal changes

Long-term measurements of radon levels in the soil shows a high variability in concentration depending on the daily changes in atmospheric pressure. It was also found that typical annual radon activity concentration in the soil air was disturbed by mild winter and heavy summer precipitation (Müllerova [2014](#)). Vasilyev & Zhukovsky ([2013](#)) confirmed the seasonal variations of radon concentrations. In a study by Moreno ([2016](#)) it was found that radon measurements presented a wide range of values. The highest soil radon levels in the vicinity of the Amer fault were found close to the fractured areas and showed very important fluctuations repeated every year, with values in summer much higher than in winter. Indoor radon measurements were taken during and after Superstorm Sandy on the US East Coast, and from the normal level of  $70 \text{ Bq m}^{-3}$  it increased up to  $1500 \text{ Bq m}^{-3}$  during the storm. The outdoor radon concentration was not significantly affected (Kotrappa [2013](#)).

The season of radon variation seems to be different in different countries, possibly due to the nature of the soil. The measured soil gas radon activity concentration around fault lines near the

North Anatolian Fault Zone in Turkey shows seasonal variation in a highly permeable sandy-gravelly soil with definite seasons without obvious long transitional periods. The summer is characterised by 1.8 times higher average soil gas radon activity concentration than the winter (Yakut [2016](#)).

Indoor radon in houses in Slovakia, Hungary & Poland shows the typical radon behaviour, with a minimum in the summer and a maximum in the winter season, whereas in 32% of schools the maximum indoor radon was reached in the summer months (Müllerová [2016](#)). This could possibly be due to ventilation?

The following conclusions were drawn as a result of a study on vegetable growing circumstances by Li ([2016](#)): Firstly, the average radon levels in typical months in Shouguang county are all much higher than that in ordinary dwellings in China, diurnal and seasonal variations in radon levels are observed inside greenhouses. Secondly, temperature and relative humidity may play a role indirectly through affecting soil moisture and other factors.

## Water supplies


A **Turkish** study showed that some spa waters contained radon at higher levels than our action levels. If people bathe indoors in such spa waters the levels of radon they inhale could add significantly to their exposure (Oner [2013](#), Akkaya [2016](#)). Radon-contaminated drinking water from private wells in the **USA** was felt to present a significant public health concern (Cappello [2013](#)). Significantly high levels of radon gas were found in **Iranian** water (Pirsaheb [2013](#)). Yang reported ([2014](#)) that wells within 5 km of granitic intrusions are at risk of containing high levels of radon.

At least 10% of people living in the **Czech Republic** are supplied with water from private sources (wells and boreholes). The official guideline limit does not apply to private sources of drinking water. Radon in water influences human health by ingestion and also by inhalation when radon is released from water during showering and cooking (Otahal [2014](#), Moldovan [2014](#)).

## Health Effects

Between 3,540 and 6,510 new cancer cases in Ontario each year result from environmental factors, says a report in August 2016 from Cancer Care Ontario and Public Health Ontario (PHO). Dr Ray Copes, chief of environmental and occupational health for PHO says "Interestingly, three main carcinogens stand out as being particularly important -- ultraviolet (UV) radiation from the sun, radon gas, and fine particle (PM2.5) air pollution." The report finds that 1,080 to 1,550 new cancer cases are associated with radon exposure. See

[http://www.publichealthontario.ca/en/eRepository/Environmental\\_Burden\\_of\\_Cancer\\_in\\_Ontario\\_2016.pdf](http://www.publichealthontario.ca/en/eRepository/Environmental_Burden_of_Cancer_in_Ontario_2016.pdf)



Radon and its progeny have been confirmed as being a category I carcinogenic agent. Data from a study by Liu (2016) demonstrated that p53-associated metabolic pathways may be altered in radon-mediated malignant transformation.

Radon changed a total of 208 genes in lung cells, 32% upregulated and 68% downregulated and depended on the dose. Further analysis showed them to be involved in biological processes related to cell cycle control/mitosis, chromosome instability and cell differentiation (Chauhan & Howland 2014).

Chen (2013) found that among those people living in homes with very high radon concentrations, it is typically parents of young children that demonstrate a great deal of concern. They want to know the risk of developing lung cancer when a child has lived in a home with high radon for a few years. The results of the study demonstrated clearly that the higher the radon concentration, the sooner remedial measures should be undertaken. There is a critical need to raise parental awareness about child health related to home exposure to radon and secondhand smoke (Huntington-Moskos 2016). Public health nurses can create promote home testing for radon and low-cost resources to reduce risk.

A study by Larionov (2016) suggested that particular types of DNA damage found in children with long-term exposure to radon could be linked to a sensitivity to radiation.

The highest rogue cell (RC) frequency was present in children environmentally exposed to radon. In 85% of RCs, double minutes, observed in a large number of human tumours, were present (Druzhinin 2016).

Exposure to radon is the second leading cause of lung cancer, especially for women in developed countries (Serke 2013), and the risk is significantly higher for smokers than for nonsmokers. Bräuner (2012) found a positive association between radon and lung cancer risk. Lung cancer in those who have never smoked accounts for between 9 and 28% of all lung cancer cases (Hubaux 2012, Truta 2014). Exposure to radon can lead to genetic and epigenetic alterations in tumour genomes, impacting genes and pathways involved in lung cancer development.

A study by Bräuner (2013) found significant associations and exposure-response patterns between long-term residential radon exposure in a general population and risk of primary brain tumours. Another study (Bräuner 2015) found that long-term residential radon exposure may contribute to development of basal cell carcinoma of the skin, though radon levels were calculated rather than measured and more follow-up is recommended.

In Spain, 14% of Galician municipalities had radon concentrations above the United States Environmental Protection Agency (USEPA) action level. There was a significant correlation between residential radon and oesophageal cancer mortality for males (Ruano-Ravina 2014).

Bräuner (2010) suggested that air pollution from traffic might enhance the effect of radon on the risk of childhood leukaemia, and Pedersen (2014) suggested that distance to powerlines together with domestic radon levels may impact the risk of childhood leukaemia. Schwartz & Klug (2015) found that incidence rates for chronic lymphocytic leukaemia (CLL) were significantly correlated with residential radon levels. Laurent (2013) concluded that there could be a link between radon, ionising radiation and leukaemia in French children, though further research was needed due to some uncertainty about the level of risk. A large, prospective study suggest residential radon may be a risk factor for lymphoid malignancies among women but not men (Teras 2016).

A large proportion of radon-attributable lung cancer deaths in Ontario are from exposures below the Canadian guideline. Peterson (2013) felt that testing and remediation may prevent a portion of radon-related lung cancer deaths.

In the first study of residential radon and birth defects, Langlois (2016) found significant associations with cleft lip with or without cleft palate and cystic hygroma / lymphangioma. They also found non-significant associations with three skeletal defects, Down syndrome, other specified anomalies of the brain, and other specified anomalies of the bladder and urethra.

## Remediation

Lantz (2013) strongly supports public communication efforts that promote residential radon testing and remediation. A questionnaire study of 23 European countries found that approximately 26,000 homes have been remediated, but millions have not yet been done. The number is increasing due to the rare use of radon prevention.

A study by the Health Protection Agency (Howarth 2013) determined the long term effectiveness of commonly used radon remedial methods over 15 years. The overall failure rate was 63%, with roughly equal rates for passive and active type systems. It was also found that all types of remedial measure can last more than 10 years, but also found examples for all measures that failed in less than 5 years. The HPA advises that homes should be retested every 5-10 years.

## Measuring radon levels in your home, school and workplace

Measuring radon levels has never been easier than with the Canary radon meter, which can give you the information you need to find out if you and your family are at risk from this invisible gas.

Kitto (2014) measured radon levels in 186 schools in New York State. Some rooms in the schools had more than 740 Bq m<sup>-3</sup>. Short-term radon measurements in the schools showed little correlation to basement and first-floor radon results from single-family homes in the towns.

Radon was measured in three nurseries and a primary school in a rural area with nongranite soil in north **Portugal**. Radon concentrations surpassed by severalfold the recommended guidelines and thresholds, and excessive levels of health concern were sporadically found (Sousa 2015).

Radon concentrations were measured in 45 classrooms from 13 public primary schools located in Porto, **Portugal**. In all schools, radon concentrations ranged from 56 to 889 Bq m<sup>-3</sup>. The results showed that the limit of 100 Bq m<sup>-3</sup> established by WHO IAQ guidelines was exceeded in 92 % of the measurements, as well as 8 % of the measurements exceeded the limit of 400 Bq m<sup>-3</sup> established by the national legislation (Madureira [2015](#)). The geometric mean of radon activity of 91 dwellings located at the Vila Pouca de Aguiar pluton in northern **Portugal** is 568Bqm<sup>-3</sup>, exceeding that of other northern Portuguese granites (Martins [2016](#)). Measurements carried out during a winter season, indicate that 62.6% of the analysed dwellings yield higher indoor radon average values than the Portuguese legislation limit of 400Bqm<sup>-3</sup>.

In British Columbia, **Canada**, some homes which lie outside of the radon-prone areas have radon levels above the Canadian guideline, which is the reason Health Canada recommends that all homes should be tested (Cheng [2016](#)).

**Hungary, Poland and Slovakia** have higher radon levels than the world average. Radon concentrations are more than 200 Bq m<sup>-3</sup> in about 87% of cases. Homes with radon concentration of about 800 Bq m<sup>-3</sup> were found in Poland and Slovakia (Müllerova [2014](#)). In **Turkey**, indoor radon concentrations were found between 1 and 1400 Bq m<sup>-3</sup> (Celebi [2015](#)).

In **Albania**, where an initial 10% of homes were measured, the indoor radon concentrations ranged from 14 to 1238 Bq m<sup>-3</sup> (Tushe [2015](#)).

All it takes is 24 hours per room to put your mind at rest, or to establish that remediation measures are needed, or old measures taken are no longer effective. EMFields **hires or sells** the radon Canary meter in the UK, and sells the meter all over the world. <http://www.emfields-solutions.com/detectors/canary.asp> You can buy online, or phone during working hours - 01353 778814.

## References

- Afolabi OT et al 2015 - *Radon level in a Nigerian University Campus* BMC Res Notes Nov 14;8:677 PMID: 26578086
- Aït Ziâne M et al 2014 - *Exposure to indoor radon and natural gamma radiation in some workplaces at Algiers Algeria* Radiat Prot Dosimetry 160(1-3):128-33 PMID: 24711531
- Akbari K et al 2013 - *Influence of indoor air conditions on radon concentration in a detached house* J Environ Radioact 116:166-73 PMID: 23159846
- Akkaya G et al 2016 - *Variation in the radon concentrations and outdoor gamma radiation levels in relation to different geological formations in the thermal regions of Bursa, Turkey* J Radiol Prot 36(3):490-503 PMID: 27355348
- Alharbi SH & RA Akber 2015 - *Radon and thoron concentrations in public workplaces in Brisbane, Australia* J Environ Radioact 144:69-76 PMID: 25827573
- Alvarez-Gallego M et al 2015 - *High radon levels in subterranean environments: monitoring and technical criteria to ensure human safety (case of Castañar cave, Spain)* J Environ Radioact 145:19-29 PMID: 25863322
- Axelsson G et al 2015 - *Lung cancer risk from radon exposure in dwellings in Sweden: how many cases can be prevented if radon levels are lowered?* Cancer causes Control 26(4):541-7 PMID: 25677843
- Aykamis AS et al 2013 - *Natural radioactivity, radon exhalation rates and indoor radon concentration of some granite samples used a construction material in Turkey* Radiat Prot Dosimetry 157(1):105-11 PMID: 23633647
- Barros N et al 2015 - *Comparative survey of outdoor, residential and workplace radon concentrations* Radiat Prot Dosimetry 163(3):325-32 PMID: 24936021
- Bochicchio F 2014 - *Protection from radon exposure at home and at work in the Directive 2013/59/Euratom* Radiat Prot Dosimetry 160(1-3):8-13 PMID: 24729590

- Borgoni R** et al 2014 – *Hierarchical modelling of indoor radon concentration: how much do geology and building factors matter?* J Environ Radioact 138C:227-237 PMID: 25261869
- Bossew P** et al 2013 – *Geographical distribution of the annual mean radon concentrations in primary schools of Southern Serbia – application of geostatistical methods* J Environ Radioact 127C:141-148 PMID: 24231373
- Botha R** et al 2016 – *Radon levels measured at a touristic thermal spa resort in Montagu (South Africa) and associated effective doses* Health Phys 111(3):281-289
- Branion-Calles MC** et al 2015 – *Evaluation of different radon guideline values based on characterization of ecological risk and visualization of lung cancer mortality trends in British Columbia, Canada* BMC Public Health Nov 19; 15(1):1144 PMID: 26584618
- Bräuner EV** et al 2015 – *residential radon exposure and skin cancer incidence in a prospective Danish cohort* PLoS One 10(8):e0135642
- Bräuner EV** et al 2013 - *Residential radon and brain tumour incidence in a Danish cohort* PLoS One 8(9):e74435 PMID: 24066143
- Bräuner EV** et al 2012 - *Residential radon and lung cancer incidence in a Danish cohort* Environ Res 118:130-6 PMID: 22749110
- Bräuner EV** et al 2010 - *Is there any interaction between domestic radon exposure and air pollution from traffic in relation to childhood leukemia risk?* Cancer Causes Control 21(11):1961-4 PMID: 20607382
- Briestensky M** et al 2014 – *Radon, carbon dioxide and fault displacements in central Europe related to the Tōhoku earthquake* Radiat Prot Dosimetry 160(1-3):78-82 PMID: 24711525
- Capello MA** et al 2013 - *Radon-contaminated drinking water from private wells: an environmental health assessment examining a rural Colorado mountain community's exposure* J Environ Health 76(4):18-24 PMID: 24341157
- Carrigan CR** et al 2016 – *Delayed signatures of underground nuclear explosions* Sci Rep Mar 16;6:23032 PMID: 26979288
- Casey JA** et al 2015 – *Predictors of indoor radon concentrations in Pennsylvania, 1989-2013* Environ Health Perspect 123(11):1130-7 PMID: 25856050
- Celebi N** et al 2015 – *Indoor radon measurements in Turkey dwellings* Radiat Prot Dosimetry 167(4):626-32 PMID: 25389360
- Chauhan V & M Howland** 2014 – *Gene Expression Responses in Human Lung Fibroblasts Exposed to Alpha Particle Radiation* Toxicol In Vitro 28(7):1222-9 PMID: 24945610
- Chen J** et al 2014 – *Results of simultaneous radon and thoron measurements in 33 metropolitan areas of Canada* Radiat Prot Dosimetry 163(2):210-6 PMID: 24748485
- Chen J** 2013 - *Canadian lung cancer relative risk from radon exposure for short periods in childhood compared to a lifetime* Int J Environ Res Public Health 10(5):1916-26 PMID: 23698696
- Cheng W** 2016 – *Radon risk communication strategies: A regional story* J Environ health 78(6):102-6 PMID: 26867298
- Choi H & P Mazzone** 2014 – *Radon and lung cancer: assessing and mitigating the risk* Cleve Clin J Med 81(9):567-75 PMID: 25183848
- Choi JR** et al 2016 – *Gene mutation discovery research of non-smoking lung cancer patients due to indoor radon exposure* Ann Occup Environ Med 28:13 PMID: 26985396
- Cosma C** et al 2015 – *Testing radon mitigation techniques in a pilot house from Băița-Ștei radon prone area (Romania)* J Environ Radioact 140:141-7 PMID: 25483354
- Dowdall A** et al 2016 – *The rate of radon remediation in Ireland 2011-2015: Establishing a base line rate for Ireland's National Radon Control Strategy* J Environ Radioact 162-163:107-112 PMID: 27227562
- Druzhinin V** et al 2016 – *Lymphocytes with multiple chromosomal damages in a large cohort of West Siberia residents: Results of long-term monitoring* Mutat Res 784-785:1-7 PMID: 26731314
- Duan P** et al 2015 – *Nonlinear dose-response relationship between radon exposure and the risk of lung cancer: evidence from a meta-analysis of published observational studies* Eur J Cancer Prev 24(4):267-77 PMID: 25117725

- Edwards JK** et al 2014 – *Occupational Radon Exposure and Lung Cancer Mortality: Estimating Intervention Effects Using the Parametric g-Formula* *Epidemiology* 25(6):829-34 PMID: 25192403
- El-Kameesy SU** et al 2016 – *Radiological safety assessment inside ancient Egyptian tombs in Saqqara* *Isotopes Environ Health Stud* 52(6):567-76 PMID: 26988604
- Fan D** et al 2016 – *Occupational exposure to radon in different kinds of non-uranium mines* *Radiat Prot Dosimetry* 170(1-4):311-4 PMID: 26940440
- Hinojosa de la Garza OR** et al 2014 – *Lung cancer mortality and radon concentration in a chronically exposed neighbourhood in Chihuahua, Mexico: a geospatial analysis* *ScientificWorldJournal* 2014:935380 PMID: 25165752
- Holmgren O** et al 2013 – *Radon remediation and prevention status in 23 European countries* *Radiat Prot Dosimetry* 157(3):392-6 PMID: 23771958
- Howarth C** 2013 - *Long term durability of radon remedial measures in a sample of UK homes* *J Radiol Prot* 33(4):763-771 PMID: 24025515
- Hubaux R** et al 2012 - *Arsenic, asbestos and radon: emerging players in lung tumorigenesis* *Environ Health Nov* 22;11:89 PMID: 23173984
- Hunter N** et al 2015 – *Calculation of lifetime lung cancer risks associated with radon exposure, based on various models and exposure scenarios* *J Radiol Prot* 35(3):539-55 PMID: 26083042
- Huntington-Moskos L** et al 2016 – *Radon, secondhand smoke, and children in the home: Creating a teachable moment for lung cancer prevention* *Public Health Nurs Jul* 22 [Epub ahead of print] PMID: 27443982
- İçhedef M** et al 2014 - *Two year evolution of radon emission and tectonic movements in Tuzla Fault Seferihisar-İzmir* 3;86C:102-108 PMID: 24530978
- Jiránek M & V Kačmařikova** 2014 - *Dealing with the increased radon concentration in thermally retrofitted buildings* *Radiat Prot Dosimetry* 160(1-3):43-7 PMID: 24729563
- Kang DR** et al – *Trends in research on indoor radon exposure and lung cancer in South Korea* *Ann Occup Environ Med Mar* 8;28:10 PMID: 26962457
- Kayakökü H** et al 2016 – *Measurements of radioactivity and dose assessments in some building materials in Bitlis, Turkey* *Appl Radiat Isot* 115:172-179 PMID: 27389882
- Keith S** et al 2012 - *Toxicological Profile for Radon* Agency for Toxic Substances and Disease Registry (US) PMID: 24049860
- Kendall GM** et al 2016 – *Variation with socioeconomic status of indoor radon levels in Great Britain: The less affluent have less radon* *J Environ Radioact* 164:84-90 PMID: 27442258
- Kim SH** et al 2016 – *Attributable risk of lung cancer deaths due to indoor radon exposure* *Ann Occup Environ Med Feb* 26;28:8 PMID: 26925236
- Kitto M** 2014 – *radon testing in schools in New York State: a 20-year summary* *J Environ Radioact* 137:213-6 PMID: 25117690
- Kotrappa P** et al 2013 - *Measurement of indoor and outdoor radon concentrations during Superstorm Sandy* *Radiat Prot Dosimetry* 157(3):455-8 PMID: 23798707
- Kourtidis K** et al 2015 – *Radon and radioactivity at a town overlying Uranium ores in northern Greece* *J Environ Radioact* 150:220-7 PMID: 26372739
- Kropat G** et al 2013 - *Major influencing factors of indoor radon concentrations in Switzerland* *J Environ Radioact* PMID: 24333637
- Kumar A** et al 2013 - *Modeling of indoor radon concentration from radon exhalation rates of building materials and validation through measurements* *J Environ Radioact* 127C:50-55 PMID: 24158045
- Langlois PH** et al 2016 – *Residential radon and birth defects: A population-based assessment* *Birth Defects Res A Clin Mol Teratol* 106(1):5-15 PMID: 25846606
- Lantz PM** et al 2013 - *Radon, smoking, and lung cancer: the need to refocus radon control policy* *Am J Public Health* 103(3):443-7 PMID: 23327258

- Larionov AV** et al 2016 – *DNA excision repair and double-strand break repair gene polymorphisms and the level of chromosome aberration in children with long-term exposure to radon* Int J Radiat Biol 92(8):466-74 PMID: 27285066
- Laurent O** et al 2013 – *Potential impacts of radon, terrestrial gamma and cosmic rays on childhood leukemia in France: a quantitative risk assessment* Radiat Environ Biophys 52(2):195-209 PMID: 23529777
- Laurier D & D Gay** 2015 – *[Risks associated to ionizing radiation from natural sources]* Rev Prat 65(1):74-8 PMID: 25842437
- Li X** et al 2016 – *Preliminary study on the variation of radon-222 inside greenhouse of Shouguang county, China* J Environ Radioact 153:120-125 PMID: 26771243
- Liu X** et al 2016 – *Radon-induced alterations in p53-mediated energy metabolism of malignantly transformed human bronchial epithelial cells* J Toxicol Environ Health A 79(9-10):436-41 PMID: 27267826
- Madureira J** et al 2015 – *Radon in indoor air of primary schools: determinant factors, their variability and effective dose* Environ Geochem Health 38(2):523-33 PMID: 26100326
- Martins LM** et al 2016 – *Indoor radon risk associated to post-tectonic biotite granites from Vila Pouca de Aguiar pluton, northern Portugal* Ecotoxicol Environ Saf 133:164-175 PMID: 27448957
- Martín Sánchez A** et al 2013 - *Additional contamination when radon is in excess* Appl Radiat Isot 81:212-5 PMID: 23548693
- Melloni BB** 2014 – *Lung cancer in never-smokers: radon exposure and environmental tobacco smoke* Eur Respir J 44(4):850-2 PMID: 25271226
- Mohery M** et al 2013 – *Radon decay products in realistic living rooms and their activity distributions in human respiratory system* Radiat Prot Dosimetry 162(3):394-9 PMID: 24324251
- Moldovan M** et al 2014 – *Radon and radium concentration in water from North-West of Romania and the estimated doses* Radiat Prot Dosimetry 162(1-2):96-100 PMID: 25031036
- Moreno V** et al 2016 – *Soil radon dynamics in the Amer fault zone: An example of very high seasonal variations* J Environ Radioact 151 Pt 1:293-303 PMID: 26551588
- Müllerová M** et al 2016 – *Indoor radon survey in Visegrad countries* Appl Radiat Isot 110:124-128 PMID: 26774389
- Müllerova M** et al 2014 – *Preliminary results of indoor radon survey in V4 countries* Radiat Prot Dosimetry 160(1-3):210-3 PMID: 24723197
- Müllerova M** et al 2014 – *Daily and seasonal variations in radon activity concentration in the soil air* Radiat Prot Dosimetry 160(1-3):222-5 PMID: 24714111
- Nafezi G** et al 2014 - *Radon levels and doses in dwellings in two villages in Kosovo, affected by depleted uranium* Radiat Prot Dosimetry 158(3):331-9 PMID: 24051175
- Nursan C** et al 2014 - *Parent's knowledge and perceptions of the health effects of environmental hazards in Sakarya, Turkey* J Pak Med Assoc 64(1):38-41 PMID: 24605711
- Oh SS** et al 2016 – *Radon exposure and lung cancer: risk in nonsmokers among cohort studies* Ann Occup Environ Med Mar 9;28:11 PMID: 26962458
- Olzowski J** et al 2015 – *[Occurrence of radon in the Polish underground tourist routes]* Med Pr 66(4):557-63 PMID: 26536972
- Oner F** et al 2013 – *Measurements of radon concentrations in spa waters in Amasya, Turkey* Radiat Prot Dosimetry 157(2):221-4 PMID: 23685534
- Otahal P** et al 2014 – *Radon in private drinking water wells* Radiat Prot Dosimetry 160(1-3):235-8 PMID: 24714110
- Pedersen C** et al 2014 – *Distance to High-Voltage Power Lines and Risk of Childhood Leukemia – an Analysis of Confounding by and Interaction with Other Potential Risk Factors* PLoS One 9(9):e107096 PMID: 25259740
- Peterson E** et al 2013 - *Lung cancer risk from radon in Ontario, Canada: how many lung cancers can we prevent?* Cancer Causes Control 24(11):2013-20 PMID: 23982909

- Pirsaheb M** et al 2013 - *A systematic review of radon investigations related to public exposure in iran* Iran Red Crescent Med J 15(11):e10204 PMID: 24719680
- Rapaglia J** et al 2015 - *A GIS typology to locate sites of submarine groundwater discharge* J Environ Radiat 145:10-18 PMID: 25863321
- Rauch SA & SB Henderson** 2013 - *A comparison of two methods for ecologic classification of radon exposure in British Columbia: residential observations and the radon potential map of Canada* Can J Public Health 104(3):e240-5 PMID: 23823889
- Rossetti M & M Esposito** 2015 - *Radon levels in underground workplaces: a map of the Italian regions* Radiat Prot Dosimetry 164(3):392-7 PMID: 25183837
- Ruano-Ravina A** et al 2014 - *Residential radon exposure and esophageal cancer. An ecological study from an area with high indoor radon concentration (Galicia, Spain)* Int J Radiat Biol 90(4):299-305 PMID: 24460133
- Schram-Bijkerk D** et al 2013 - *The burden of disease related to indoor air in the Netherlands: do different methods lead to different results?* Occup Environ Med 70(2):126-32 PMID: 23204013
- Schwartz GG & MG Klug** 2016 - *Incidence rates of chronic lymphocytic leukemia in US states are associated with residential radon levels* Future Oncol 12(2):165-74 PMID: 26515766
- Serke M** et al 2013 - *[Gender-specific difference in lung cancer]* Pneumologie 67(5):270-9 PMID: 23677552
- Signorelli C & RM Limina** 2002 - *[Environmental risk factors and epidemiologic study]* Ann Ig 14(3):253-62 PMID: 12162123
- Singh P** et al 2016 - *Theoretical modelling of indoor radon concentration and its validation through measurements in South-East Haryana, India* J Environ Manage 171:35-41 PMID: 26874612
- Smyth R** et al 2016 - *Radon testing in rapid access lung clinics: an opportunity for secondary prevention* Ir J Med Sci Mar 30 [Epub ahead of print] PMID: 27083463
- Sousa SI** et al 2015 - *Radon levels in nurseries and primary schools in Bragança District - Preliminary assessment* J Toxicol Environ Health 78(13-14):805-13 PMID: 26167747
- Stojanovska Z** et al 2016 - *Variation of indoor radon concentration and ambient dose equivalent rate in different outdoor and indoor environments* Radiat Environ Biophys 55(2):171-83 PMID: 26943159
- Tait DR** et al 2013 - *Enrichment of radon and carbon dioxide in the open atmosphere of an Australian coal seam gas field* Environ Sci Technol 47(7):3099-104 PMID: 23444905
- Tamari Y** et al 2016 - *A report that Fukushima residents are concerned about radiation from Land, Food and Radon* J Radiat Res 57(4):418-21 PMID: 26983979
- Teras LR** et al 2016 - *residential radon exposure and risk of incident hematologic malignancies in the Cancer Prevention Study-II Nutrition cohort* Environ Res 148:46-54 PMID: 27015563
- Torres-Durán M** et al 2015 - *Small cell lung cancer in never-smokers* Eur Respir J 47(3):947-53 PMID: 26699724
- Torres-Durán M** et al 2013 - *Residential radon and lung cancer in never smokers. A systematic review* Cancer Lett 345(1):21-6 PMID: 24333737
- Truta LA** et al 2014 - *Lung cancer risk due to residential radon exposures: estimation and prevention* Radiat Prot Dosimetry 160(1-3):112-6 PMID: 24751985
- Tushe KB** et al 2015 - *First step towards the geographical distribution of indoor radon in dwellings in Albania* Radiat Prot Dosimetry Dec 8 [Epub ahead of print] PMID: 26656073
- Valmari T** et al 2014 - *Radon measurement and mitigation activity in Finland* Radiat Prot Dosimetry 160(1-3):18-21 PMID: 24729562
- Vasilyev A & I Yarmoshenko** 2016 - *Effect of energy-efficient measures in building construction on indoor radon in Russia* Radiat Prot Dosimetry Jun 29 [Epub ahead of print] PMID:27358394
- Vasilyev AV & MV Zhukovsky** 2013 - *Determination of mechanisms and parameters which affect radon entry into a room* J Environ Radioact Jun 26;124C:185-90 PMID: 23811128

**Yakut H** et al 2016 – *Soil gas radon measurement around fault lines on the western section of the north Anatolian fault zone in Turkey* Radiat Prot Dosimetry Feb 21 [Epub ahead of print] PMID: 26903173

**Yang Q** et al 2014 - *Uranium and Radon in Private Bedrock Well Water in Maine: Geospatial Analysis at Two Scales* Environ Sci Technol 48(8):4298-306 PMID: 24655434

**Yarmoshenko IV** et al 2014 – *Indoor radon problem in energy efficient multi-storey buildings* Radiat Prot Dosimetry 160(1-3):53-6 PMID: 24723188

**Zhou X** et al 2015 – *Environmental impact of CO<sub>2</sub>, Rn, Hg degassing from the rupture zones produced by Wenchuan M s 8.0 earthquake in western Sichuan, China* Environ Geochem Health 38(5):1067-82 PMID: 26486131

**Zhukovsky MV & AV Vasilyev** 2014 – *Mechanisms and sources of radon entry in buildings constructed with modern technologies* Radiat Prot Dosimetry 160(1-3):48-52 PMID: 24729591

**Zunić ZS** et al 2014 – *High variability of indoor radon concentrations in uraniferous bedrock areas in the Balkan region* Appl Radiat Isot 94:328-37 PMID: 25305525